

Dry (Frequent Fire) Mixed Conifer Forest Desired Conditions

Operational Draft: This document is prepared to provide guidance to Forest Plan revision teams. As this guidance is implemented, we expect to learn improved ways to do this work. As we learn, this document will be updated. This document was reviewed and revised as appropriate in April 2010 to conform to the requirements of the 1982 Planning Rule provisions.

General Description

The dry mixed conifer forest vegetation community is transitional with increasing elevation between ponderosa pine and wet mixed-conifer forests and generally occurs at elevations ranging from approximately 5,500 to 9,500 feet. Dry mixed-conifer forests are dominated by mainly shade intolerant trees such as ponderosa pine, southwestern white pine, limber pine, quaking aspen, and Gambel oak, with a lesser presence of shade tolerant species such as white fir and blue spruce. Mid-tolerant species such as Douglas-fir are common. Aspen may occur as individual trees or small groups. This forest vegetation community typically occurs with an understory of grasses, forbs, and shrubs.

Landscape Scale Desired Conditions (10,000 + acres)

At the landscape scale, the dry mixed conifer vegetation community is a mosaic of forest conditions composed of structural stages ranging from young to old trees. Forest appearance is variable but generally uneven-aged and open; occasional patches of even-aged structure are present. The forest arrangement is in small clumps and groups of trees interspersed within variably-sized openings of grass/forb/shrub vegetation associations similar to historic patterns. Openings typically range from 10 percent in more productive sites to 50 percent in the less productive sites. Size, shape, number of trees per group, and number of groups per area are variable across the landscape. Where they naturally occur, groups of aspen and all structural stages of oak are present. Denser tree conditions exist in some locations such as north facing slopes and canyon bottoms.

Old growth occurs throughout the landscape, generally in small areas as individual old growth components, or as clumps of old growth. Old growth components include old trees, dead trees (snags), downed wood (coarse woody debris) and structural diversity. The location of old growth shifts on the landscape over time as a result of succession and disturbance (tree growth and mortality).

The dry mixed conifer forest vegetation community is composed predominantly of vigorous trees, but declining trees are a component and provide for snags, top-killed, lightning- and fire-scarred trees, and coarse woody debris (>3 inch diameter), all well-distributed throughout the landscape. Snags are typically 18 inches or greater at DBH and average 3 per acre. Downed logs (>12 inch diameter at mid-point, >8 feet long) average 3 per acre within the forested area of the landscape. Coarse woody debris, including downed logs, ranges from 5 to 15 tons per acre.

The composition, structure, and function of vegetative conditions are resilient to the frequency, extent, severity of disturbances, and to climate variability. The landscape is a functioning ecosystem that contains all its components, processes, and conditions that result from endemic levels of disturbances (e.g. insects, diseases, fire, and wind), including snags, downed logs, and old trees. Grasses, forbs, shrubs, needle cast (fine fuels), and small trees maintain the natural fire regime. Organic ground cover and herbaceous vegetation provide protection of soil, moisture infiltration, and contribute to plant and animal diversity and to ecosystem function. Frequent, low severity fires (Fire Regime I) are characteristic, including throughout goshawk home ranges. Natural and anthropogenic disturbances are sufficient to maintain desired overall tree density, structure, species composition, coarse woody debris, and nutrient cycling.

Mid-Scale Desired Conditions (100 -1,000 acres)

At the mid-scale the dry mixed conifer forest vegetation community is characterized by variation in the size and number of tree groups depending on elevation, soil type, aspect, and site productivity. The more biologically productive sites contain more trees per group and more groups per area. Openings typically range from 10 percent in more productive sites to 50 percent in the less productive sites. Tree density within forested areas generally ranges from 30 to 100 square foot basal area per acre.

The mosaic of tree groups generally comprises an uneven-aged forest with all age classes and structural stages. Occasionally small patches (generally less than 50 acres) of even-aged forest structure are present. Disturbances sustain the overall age and structural distribution.

Fires burn primarily on the forest floor and do not spread between tree groups as crown fire.

Forest structure in the wildland urban interface (WUI)¹ has smaller and more widely spaced groups of trees than in the non-WUI areas.

Forest conditions in goshawk post-fledging family areas (PFAs) are similar to general forest conditions except these forests contain 10 to 20 percent higher basal area in mid-aged to old tree groups than in goshawk foraging areas and in the general forest. Goshawk nest areas have forest conditions that are multi-aged but are dominated by large trees with relatively denser canopies than other areas in the dry mixed conifer type.

Fine Scale Desired Conditions (≤ 10acres)

Trees typically occur in irregularly shaped groups and are variably-spaced with some tight clumps. Crowns of trees within the mid-aged to old groups are interlocking or nearly interlocking. Openings surrounding tree groups are variably-shaped and comprised of a grass/forb/shrub mix. Some openings contain individual trees or snags.

¹ Note –each Forest needs to provide the definition for WUI that they are using.

Trees within groups are of similar or variable ages and one or more species. Size of tree groups typically is less than 1 acre. Groups at the mid-age to old stages consist of 2 to approximately 50 trees per group.