

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG)

R4OKHK Oak Woodland

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

John Pearson John.Pearson@dnr.state.ia.us

Reviewers

John Ortmann jortmann@tnc.org
George Hartman george.hartman@mdc.mo.gov
Jim Drake jim_drake@natureserve.org

Vegetation Type

Woodland

General Model Sources

- Literature
 Local Data
 Expert Estimate

Rapid Assessment Model Zones

- California Pacific Northwest
 Great Basin South Central
 Great Lakes Southeast
 Northeast S. Appalachians
 Northern Plains Southwest
 N-Cent.Rockies

Dominant Species*

QUAL TIAM
QURU CAOVS
QUMA OSVI
ACSA QUMU

LANDFIRE Mapping Zones

43
42

Geographic Range

Occurring primarily in Iowa, northern Missouri, eastern Nebraska.

Biophysical Site Description

Found in wooded valleys extending into the extensive landscape of tallgrass prairie on the uplands. Topographically, distributed on dry and dry-mesic slopes between flat uplands and bottomlands; also on dry sites on flat uplands adjacent to edges of valleys. Generally, from east to west, the distribution becomes more and more limited in extent and more dependent on favorable habitat conditions. Open conditions describe a single canopy structure with no developed midstory. Closed conditions are multiple canopy usually late-seral forests.

Vegetation Description

White oak, red oak, bur oak, basswood, sugar maple (east), and black maple (west) are the main tree species; bur oak, hickory, white ash, and American and red elms are frequent associates. Ironwood and roughleaf dogwood are important under and mid story components. Toward the western edge of the type, the following tree species drop out of the model due to geographic range: sugar maple, black maple, white, oak, red oak. At the far western edge, only bur oak and basswood remain as major canopy trees.

Disturbance Description

Fire Regime Group I. Frequent surface fires (5-yr fire return interval) in the understory under woodland density of canopy trees. Interruptions to frequent fire are necessary to allow tree regeneration to replace aging canopy trees. A 15-yr pause in burning is modeled as opportunity for regenerating trees to grow into fire-resistant size class before surface fire resumes.

Adjacency or Identification Concerns

Adjoins oak savanna (R4OASA) toward prairie side of woodland. Adjoins maple-basswood forest communities on rarely burned mesic sites. Model focuses on portion of landscape that could support oak

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

woodland.

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

This PNVG was found in fairly protected areas with minimum patch sizes 100-1000 and maximum 10,000 acres.

Issues/Problems

Class A (open regenerating state) modeled out to be 16% of outcome, but presettlement condition was likely much less (1%).

Model Evolution and Comments

Ortmann suggested that grazing/browsing could have been an important influence in establishment of canopy trees in gaps and in recruitment/establishment of understory species.

Succession Classes**
Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 16 %

Early1 All Struct

Description

0-15years. Sprouts, seedlings, saplings of major overstory species in gaps and openings created by wind, insect/disease and fire. Shrubs abundant as well. Both fire-tolerant and intolerant species present. Trees grow into fire-resistant sizes by end of this class.

Dominant Species* and Canopy Position

QUAL Upper
QURU Upper
TIAM Upper
ACSA Upper

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 1

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	100 %
Height	Herb Short <0.5m	Tree Regen <5m
Tree Size Class	Sapling >4.5ft; <5"DBH	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 67 %

Late1 Open

Description

16-400 years. Single canopy structure without woody understory. Herbaceous layer dense. White oak and red oak are dominant species due to fire exclusion of maple and basswood. After 150 years, red oak dies of old age, leaving white oak as main dominant.

Dominant Species* and Canopy Position

QUAL Upper
QURU Upper

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model 1

Structure Data (for upper layer lifeform)

	Min	Max
Cover	25 %	60 %
Height	Tree Short 5-9m	Tree Medium 10-24m
Tree Size Class	Large 21-33"DBH	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class C 17%

Late1 Closed

Description

16-400 years. Mature canopy. White oak is dominant overstory species variable due to death of red oak and fire exclusion of maple and basswood.

Dominant Species* and Canopy Position

ACSA Upper
TIAM Upper
QURU Upper
QUAL Upper

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	60 %	100 %
Height	Tree Short 5-9m	Tree Medium 10-24m
Tree Size Class	Large 21-33"DBH	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 0%

Mid1 Closed

Description

Dominant Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	0 %
Height		
Tree Size Class		

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class E 0%

Late1 Closed

Description

Dominant Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	%	%
Height		
Tree Size Class		

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Disturbances

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Disturbances Modeled

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

Historical Fire Size (acres)

Avg: no data
 Min: no data
 Max: no data

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

Fire Regime Group: 1

I: 0-35 year frequency, low and mixed severity
 II: 0-35 year frequency, replacement severity
 III: 35-200 year frequency, low and mixed severity
 IV: 35-200 year frequency, replacement severity
 V: 200+ year frequency, replacement severity

Fire Intervals (FI)

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
<i>Replacement</i>	450			0.00222	2
<i>Mixed</i>		0	0		
<i>Surface</i>	7.5			0.13333	98
<i>All Fires</i>	7			0.13557	

References

Anderson, Pail F. 1996. GIS research to digitize maps of Iowa 1832-1859 vegetation from General Land Office township plat maps. Iowa Department of Natural Resources, Des Moines, Iowa. See also: <http://www.public.iastate.edu/~fridolph/dnrglo.html>

Cleveland, M.K. and D.N. Duvick. 1992. Iowa climate reconstructed from tree rings, 1640-1982. *Water Resources Research* 28(10):2607-2615.

Duvick, Daniel and T.J. Blasing. 1983. Iowa's oldest oaks. *Proceedings of the Iowa Academy of Science* 90(1):32-34.

Fralish, James S., Fred B. Crooks, Jim L. Chambers, and Francis M. Harty. 1991. Comparison of presettlement, second-growth, and old-growth forest on six site types in the Illinois Shawnee Hills. *American Midland Naturalist* 125:294-309.

Pearson, John. 1989. Ancient record keepers. *Iowa Conservationist*, February 1988.

Tester, John R. 1989. Effects of fire frequency on oak savanna in east-central Minnesota. *Bulletin of the Torrey Botanical Club* 116(2):134-144.

Thomson, George W. and H. Gene Hertel. 1981. The forest resources of Iowa in 1980. *Proceedings of the Iowa Academy of Science* 88(1):2-6.

Thomson, George W. 1987. Iowa's forest area in 1832: a reevaluation. *Proceedings of the Iowa Academy of Science* 94(4):116-120.

Van Lear, David H. and Thomas Waldrop. 1989. History, uses, and effects of fire in the Appalachians. General Technical Report SE-54, Southeastern Forest Experiment Station, USDA Forest Service. 20 pp.

White, Alan S. 1983. The effects of thirteen years of annual prescribed burning on a *Quercus ellipsoidalis* community in Minnesota. *Ecology* 64(5):1081-1085.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.