

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG)

R#DFHEwt Douglas-fir Hemlock-Wet Mesic

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Jane Kertis jkertis@fs.fed.us
Karen Kopper karen_kopper@nps.gov
Steve Acker steve_acker@nps.gov

Reviewers

Vegetation Type

Forested

Dominant Species*

PSME
TSHE
THPL

General Model Sources

- Literature
 Local Data
 Expert Estimate

LANDFIRE Mapping Zones

1 8
2 9
7

Rapid Assessment Model Zones

- California Pacific Northwest
 Great Basin South Central
 Great Lakes Southeast
 Northeast S. Appalachians
 Northern Plains Southwest
 N-Cent.Rockies

Geographic Range

This type occupies low montane elevations of western Washington and Oregon. In Washington it occurs on the north, south and west side of the Olympic Peninsula and along the low to moderate elevation western slopes of the Cascade Range. In Oregon this type is found in mesic to wet microsites, on northerly slopes and upper elevations of the Cascades, and on the west side and upper east side of the Coast Range.

Biophysical Site Description

Soils vary from well-drained to clay. This type commonly occupies mesic to wet to cool microsites on all aspect at elevations up to 4000 feet in elevation.

Vegetation Description

Douglas-fir and western hemlock dominate this PNVG. Western red cedar is a common associate. Common understory herbs and shrubs include Devil's club, various huckleberry species, vine maple, rhododendron, Oregon oxalis, bear grass, swordfern and bunchberry dogwood.

Disturbance Description

Fire plays a major role in infrequently resetting landscapes within this PNVG, with intervals ranging roughly from 300 to 800 years. Mixed severity fires occur less frequently in this regime than in the Douglas-fir Hemlock mesic dry regime.

Insects, pathogens and windthrow occur in this type at variable intervals, creating fine scale variability on the landscape.

Adjacency or Identification Concerns

The Silver fir low type bounds this PNVG at higher elevations, and the Douglas-fir Hemlock mesic dry can bound it at lower elevations and drier microsites.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

Although fires are often large (100s-1000 acres), fire severity patterns are quite variable, ranging from underburns to high severity patches within single events. Wind, insects and pathogens can create gaps of various sizes.

Issues/Problems

Model Evolution and Comments

Succession Classes**

Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 5%

Early1 PostRep

Description

Post-stand replacement community consisting of herbs, and/or shrubs such as bracken fern, fireweed, ceanothus. Douglas-fir, western hemlock and western red cedar seedlings may be present.

Dominant Species* and Canopy Position

PTERI
CHAN9
PSME
TSHE

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	10 %	100 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 15%

Mid1 Closed

Description

Closed-canopy young forest stands with trees up to 20 inches in diameter, usually conifers (especially Douglas-fir and western hemlock), but with hardwoods in some cases (e.g., bigleaf maple, red alder). Understory tends to be minimal because of low light levels.

Dominant Species* and Canopy Position

PSME
TSHE
ALRU2
ACMA

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	60 %	100 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class C 1%

Mid1 Open

Description

These are young forest stands that have been opened up by mixed-severity fire. Trees are up to 20 inches in diameter. The dominant tree species is Douglas-fir, although western hemlock may be present. Shrubs such as huckleberry, rhododendron, vine maple and Devils club dominate the understory, although herbs such

Dominant Species* and Canopy Position

PSME
GASH
VACCI
POMU

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	20 %	60 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

as Oregon oxalis, , and swordfern may have appreciable cover.

Class D 4%

Late I Open

Description

These are mature to old-growth forest stands that have been opened up by mixed-severity fire. The largest trees are greater than 20 inches in diameter. The degree of canopy opening may be sufficient to permit recruitment of shade-intolerant species (e.g., Douglas-fir), or may only permit recruitment of western hemlock and other shade-tolerant species. This class has a diverse understory with essentially the same species as class E.

Dominant Species* and Canopy Position

PSME
TSHE
GASH
POMU

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	20 %	60 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class E 75%

Late I Closed

Description

These are mature to old-growth forest stands dominated by large individuals (>20 inches in diameter) of Douglas-fir and western hemlock, with advanced regeneration of western hemlock. Understories can be a mixed of shrubs such as huckleberry and vine maple, and herbs such as Oregon oxalis, bunchberry dogwood, swordfern, and twinflower.

Dominant Species* and Canopy Position

PSME
TSHE
GASH
MANE2

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	60 %	100 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Disturbances

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Disturbances Modeled

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

Historical Fire Size (acres)

Avg: no data
 Min: no data
 Max: no data

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

Fire Regime Group: 5

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

Fire Intervals (FI)

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
<i>Replacement</i>	400			0.0025	71
<i>Mixed</i>	1000			0.001	28
<i>Surface</i>					
<i>All Fires</i>	286			0.00351	

References

Agee, J. K. 1993. Fire ecology of Pacific Northwest forests. Washington, D.C.: Island Press.

Cissel, JH, et al. 1998. A landscape plan based on historical fire regimes for a managed forest ecosystem: the Augusta Creek Study. USDA PNW Gen Tech. Rep PNW-GTR-422. 82pp.

Franklin, J. F., and C. T. Dyrness. 1973. Natural vegetation of Oregon and Washington. Corvallis OR: Oregon State University Press.

Garza, E.S. 1995. Fire History and Fire Regimes of East Humbug and Scorpion Creeks and their relation to the range of Pinus lambertiana Dougl. 75 pp.

Henderson, J.A., D.H. Peter, R.D. Leshner, and D.C. Shaw. 1989. Forested Plant Associations of the Olympic National Forest. USDA Forest Service, Pacific Northwest Region. R6 ECOL Technical Paper 001-88. 502 p.

Morrison, P. H., and F. J. Swanson. 1990. Fire history and pattern in a Cascade Range landscape. PNW-GTR-254. USDA Forest Service, Pacific Northwest Research Station.

McCain, C and N. Diaz 2002. Field Guide to the Forested Plant Associations of the Westside Central Cascades of Northwest Oregon. USDA FS PNW Tech Paper R6-NR-ECOL-TP-02-02

McCain, C and N. Diaz 2002. Field Guide to the Forested Plant Associations of the Northern Oregon Coast Range USDA FS PNW Tech Paper.

Teensma, P. D., 1987. Fire history and fire regimes of the central western cascades of Oregon. Ph.D. Dissertation, University of Oregon.

Weisberg, P.J. 1998. Fire History, Fire Regimes and Development of Forest Structure in the Central Western Oregon Cascades. PhD dissertation. Oregon State University. 256 pp

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.