

## Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit [www.landfire.gov](http://www.landfire.gov). Please direct questions to [helpdesk@landfire.gov](mailto:helpdesk@landfire.gov).

### Potential Natural Vegetation Group (PNVG)

R9PIRO Pine Rocklands

#### General Information

**Contributors** (additional contributors may be listed under "Model Evolution and Comments")

**Modelers**

Duane Weis      [weisd@doacs.state.fl.us](mailto:weisd@doacs.state.fl.us)

**Reviewers**

Chris Bergh      [cbergh@tnc.org](mailto:cbergh@tnc.org)  
Rick Anderson      [thomas\\_r\\_anderson@nps.gov](mailto:thomas_r_anderson@nps.gov)  
Hillary Cooley      [hillary\\_cooley@nps.gov](mailto:hillary_cooley@nps.gov)

**Vegetation Type**

Woodland

**General Model Sources**

- Literature
- Local Data
- Expert Estimate

**Rapid Assessment Model Zones**

- California
- Great Basin
- Great Lakes
- Northeast
- Northern Plains
- N-Cent.Rockies
- Pacific Northwest
- South Central
- Southeast
- S. Appalachians
- Southwest

**Dominant Species\***

PIELD      ANCA  
SERE2      SCRH  
THRIN      RAPU  
GUSC

**LANDFIRE Mapping Zones**

56

#### Geographic Range

Pine rocklands occur in extreme south Florida and the lower Florida Keys.

#### Biophysical Site Description

Pine rocklands occur on alkaline limestone bedrock.

#### Vegetation Description

The overstory consists primarily of south Florida slash pine (*Pinus elliotti* var. *densa*) with crown closure ranging from 10 to 60%. A sometimes sparse, but often species-rich understory consists of shrubby tropical evergreen hardwoods, palms, forbs, and graminoids. Common palms include thatch palm (*Thrinax morrisii*, *T. radiata*), silver palm (*Coccothrinax argentata*), saw palmetto (*Serenoa repens*), and cabbage palm (*Sabal palmetto*). Common shrubs or subcanopy species include live oak (*Quercus virginiana*), wild tamarind (*Lysiloma latisiliquum*), poisonwood (*Metopium toxiferum*), indigo berry (*Randia aculeata*), varnish leaf (*Dodonea viscosa*), myrsine (*Rapanea punctata*), rough velvet seed (*Guettarda scabra*) cocoplum (*Chrysobalanus icaco*), willow bustic (*Bumelia salicifolia*), and marlberry (*Ardisia escallonioides*). Typical graminoid and forb species include splitbeard bluestem (*Andropogon cabanisii*), little bluestem (*Schizachyrium rhizomatum*), showy milkwort (*Polygala grandiflora*), pineland heliotrope (*Heliotropium polyphyllum*), silver dwarf morning glory (*Evolvulus sericeus*), and rabbitbells (*Crotalaria rotundifolia*).

#### Disturbance Description

This PNVG is classified as a Fire Regime Group I, 1-5 year mean fire return interval, with frequent, low intensity fires occurring at any time of year. Most acreage burns from April to June during the early lightning season. Less common (1-2 /decade) moderately severe fires associated with drought occur primarily in March to May. Anthropogenic fire was considered but is not expected to change reference class composition.

Bergh, in his review of the model, stated that a 1-5 year mean fire return interval may be too frequent. His

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

estimate ranged from 3-10 years.

**Adjacency or Identification Concerns**

Pine rocklands are often interrupted by patches of tropical hardwood hammock, which will invade into the pinelands in the absence of fire.

**Scale Description**

Sources of Scale Data  Literature  Local Data  Expert Estimate

This PNVG occurs in patches ranging in size from 200 to 10,000 acres in areas where the soil depth is minimal due to the presence of pinnacle rock. These patches were likely fragmented by the presence of tropical hardwood stands, everglades marsh, and cypress domes or savannahs.

**Issues/Problems**

The natural fire regime is currently altered by urbanization and artificially controlled water levels. Invasive exotics include Burma reed and Brazilian pepper.

**Model Evolution and Comments**

FRCC model (SFSP1) developed by Caroline Noble for Pine Rocklands and South Florida Slash Pine was used with no changes to the VDDT model. Information in the database was edited to specifically address Pine Rocklands.

**Succession Classes\*\***  
*Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).*

**Class A 15 %**

Early1 All Struct

**Description**

Class A, 0-15 years post replacement, includes seedlings, saplings, and poles of south Florida slash pine. Individual tree gaps and clusters interspersed throughout the landscape result from mortality from wind or lightning.

**Dominant Species\* and Canopy Position**

ANCA2 Lower  
SCRH Lower  
GUSC Low-Mid  
SERE2 Low-Mid

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Structure Data (for upper layer lifeform)**

|                 | Min            | Max             |
|-----------------|----------------|-----------------|
| Cover           | 10 %           | 50 %            |
| Height          | Tree Regen <5m | Tree Short 5-9m |
| Tree Size Class | Pole 5-9" DBH  |                 |

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

The dominant life form includes grasses, forbs, and small shrubs with a canopy closure of 50 to 75% and a height of less than 0.5m.

**Fuel Model 2**

**Class B 5 %**

Mid1 Closed

**Description**

Class B occurs from 16-49 years post replacement and includes mid-story development of a shrub layer. Hardwood and palm encroachment is becoming increasingly dense. This class may be the result of mosaic hammock fire.

**Dominant Species\* and Canopy Position**

PIELD Upper  
SAPA Mid-Upper  
THRIN Middle  
SERE2 Middle

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Structure Data (for upper layer lifeform)**

|                 | Min             | Max                |
|-----------------|-----------------|--------------------|
| Cover           | 10 %            | 30 %               |
| Height          | Tree Short 5-9m | Tree Medium 10-24m |
| Tree Size Class | Pole 5-9" DBH   |                    |

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

The dominant life form begins to transition to the shrub layer, primarily saw palmetto and tropical hardwoods. Canopy closure in the shrub layer increases to 25 to 40% with an average height of 1m.

**Fuel Model 5**

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Class C 25%**

Mid1 Open  
**Description**

Class C occurs from 16-49 years post replacement. There is less than 40% tree canopy closure represented by scattered individual slash pines. The understory is comprised of grasses, forbs, low shrubs and palms.

**Dominant Species\* and Canopy Position**

PIELD Upper  
SERE2 Middle  
ANCA2 Lower  
SCRU Lower

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Structure Data (for upper layer lifeform)**

|                        | <i>Min</i>      | <i>Max</i>         |
|------------------------|-----------------|--------------------|
| <i>Cover</i>           | 10 %            | 40 %               |
| <i>Height</i>          | Tree Short 5-9m | Tree Medium 10-24m |
| <i>Tree Size Class</i> | Medium 9-21"DBH |                    |

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

The dominant lifeform remains the grasses and forbs mixed with small isolated patches of shrubs.

**Fuel Model 2**

**Class D 50%**

Late1 Open  
**Description**

Trees in Class D are 50+ years old. There is less than 30% tree canopy closure, with tree diameters up to 21" dbh. The understory is comprised of grasses, forbs, low shrubs and palms.

**Dominant Species\* and Canopy Position**

PIELD Upper  
SERE2 Middle  
ANCA2 Lower  
SCRU Lower

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Structure Data (for upper layer lifeform)**

|                        | <i>Min</i>         | <i>Max</i>         |
|------------------------|--------------------|--------------------|
| <i>Cover</i>           | 10 %               | 30 %               |
| <i>Height</i>          | Tree Medium 10-24m | Tree Medium 10-24m |
| <i>Tree Size Class</i> | Medium 9-21"DBH    |                    |

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

The dominant lifeform remains the grasses and forbs mixed with small isolated patches of shrubs.

**Fuel Model 2**

**Class E 5%**

Late1 Closed  
**Description**

Trees in Class E are 50+ years old. With continued exclusion of fire, the transition to tropical hardwood hammock will begin. The dominant species include slash pine, cabbage palm, and tropical hardwoods.

**Dominant Species\* and Canopy Position**

PIELD Upper  
SAPA Mid-Upper  
LYLA3 Mid-Upper  
QUVI Mid-Upper

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Structure Data (for upper layer lifeform)**

|                        | <i>Min</i>         | <i>Max</i>         |
|------------------------|--------------------|--------------------|
| <i>Cover</i>           | 40 %               | 60 %               |
| <i>Height</i>          | Tree Medium 10-24m | Tree Medium 10-24m |
| <i>Tree Size Class</i> | Medium 9-21"DBH    |                    |

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Disturbances**

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Disturbances Modeled**

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

**Historical Fire Size (acres)**

Avg: 1500  
 Min: 1000  
 Max: 5000

**Sources of Fire Regime Data**

- Literature
- Local Data
- Expert Estimate

**Fire Regime Group: 1**

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

**Fire Intervals (FI)**

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

|                    | <i>Avg FI</i> | <i>Min FI</i> | <i>Max FI</i> | <i>Probability</i> | <i>Percent of All Fires</i> |
|--------------------|---------------|---------------|---------------|--------------------|-----------------------------|
| <i>Replacement</i> | 800           |               |               | 0.00125            | 0                           |
| <i>Mixed</i>       | 330           |               |               | 0.00303            | 1                           |
| <i>Surface</i>     | 3             | 1             | 5             | 0.33333            | 99                          |
| <i>All Fires</i>   | 3             |               |               | 0.33761            |                             |

**References**

Brown, James K. and Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Everglades National Park Fire Management plan and Environmental Assessment (Draft 2003). Everglades National Park.

Myers, Ronald L. and Ewel, John J., eds. 1990. Ecosystems of Florida. Orlando, FL: University of Central Florida Press. 765 p.

Schmidt, Kirsten M., Menakis, James P., Hardy, Colin C., Hann, Wendel J. and Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.

U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/>.

\*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.