



5.2 Plants that attract beneficial insects

Buffers can be planted with plants to attract beneficial insects that prey on insect pests. Considerations include selecting plants that bloom sequentially throughout the growing season and avoid plants that enhance pest abundance.

Native plants are preferred because these species attract native beneficial insects and are less likely to become agricultural weeds. The table provides a list of beneficial insects, the pests the beneficial insects prey on, and the plants or habitats that attract the beneficial insects.

Plants that Attract Beneficial Insects		
Beneficial	Pests	Plants/Habitat
Assassin bug (Reduviidae family)	Many insects including flies and large caterpillars	Permanent plantings for shelter (e.g., windbreaks)
Bees-Butterflies (Many families)	None but important for pollination	Pea, borage, and aster families, milkweeds, butterfly bush, others
Braconid wasp (Braconidae family)	Armyworm, cabbageworm, codling moth, gypsy moth, European corn borer, aphid, caterpillars, and other insects	Nectar plants with small flowers, yarrow, sunflower, cowparsnip
Damsel bug (Nabidae family)	Aphid, thrips, leafhopper, treehopper, caterpillars	Aster family, yarrow, common boneset
Ground beetle (Carabidae family)	Slug, snail, cutworm, Colorado potato beetle, gypsy moth, caterpillars, weed seeds	Amaranth, bunch grasses, permanent plantings for shelter

Plants that Attract Beneficial Insects		
Beneficial	Pests	Plants/Habitat
Hover fly (Syrphidae family)	Aphid	Carrot and aster family (coreopsis, sunflowers, goldenrod), cowparsnip, common boneset
Lacewing (Chrysopidae family)	Soft bodied insects including aphid, thrips, European corn borer, mealybug, scale, mite	Carrot and aster family (coreopsis, sunflowers, goldenrod)
Ladybug beetle (Coccinellidae family)	Aphid, spider mite, European corn borer, mealybug	Aster family, butterfly weed, native grasses, giant hyssop, cowparsnip, yarrow, black locust
Minute Pirate Bug (Anthocorid family)	Thrips, spider mite, leafhopper, corn earworm, small caterpillars, and other insects	Carrot and aster family (daisies, sunflowers, yarrow, goldenrod), blue elderberry, potentilla, giant hyssop, common boneset, and willows
Rove beetle (Staphylinidae family)	Aphid, nematode, flies	Native grasses, permanent plantings for shelter
Spider (Salticidae, Thomisidae, and other families)	Many insects	Carrot and aster family, giant hyssop
Spined soldier bug (<i>Podisus maculiventris</i>)	Armyworm, sawfly, Colorado potato beetle, Mexican bean beetle	Aster family (sunflowers, yarrow)
Tachinid fly (Tachinidae family)	Cutworm, armyworm, May beetle, gypsy moth, squash bug	Carrot and aster family, amaranth
Tiger beetle (Cicindelidae family)	Many insects	Amaranth, bunch grasses, permanent plantings for shelter
Chalcid wasps (many families including Trichogrammatidae)	Spruce budworm, cotton bollworm, tomato hornworm, corn earworm, corn borer, codling moth	Carrot and aster family (daisies, sunflowers, yarrow, goldenrod), potentilla, giant hyssop, cowparsnip, common boneset

5.2 References

- Al-Doghairi, M.A.; Cranshaw, W.S.. 1999. Surveys on visitation of flowering landscape plants by common biological control agents in Colorado. *Journal of the Kansas Entomological Society*. 72: 190-196.
- Altieri, M.A.; Nicholls, C.I.; Fritz, M.A. 2005. Manage insects on your farm: a guide to ecological strategies. Beltsville, MD: Sustainable Agriculture Network. 130 p. <http://www.sare.org/publications/insect/insect.pdf>. [Date accessed: September 25, 2007].
- Bugg, R. 1994. Using cover crops to manage arthropods of orchards: A review. *Agriculture, Ecosystems and Environment*. 50:11–28.
- Colley, M.R.; Luna, J.M. 2000. Relative attractiveness of potential beneficial insectary plants to aphidophagous hoverflies. *Environmental Entomology*. 29: 1054-1059.
- Defour, R. 2000. Farmscaping to enhance biological control. Fayetteville, AR: Appropriate Technology Transfer for Rural Areas. 40 p. <http://attra.ncat.org/attra-pub/PDF/farmscaping.pdf>. [Date accessed: September 26, 2007].
- Fiedler, A.; Tuell, J.; Issacs, R.; Landis, D. 2007. Attracting beneficial insects with native flowering plants. Bulletin E-2973. East Lansing, MI: Michigan State University. 5 p. <http://nativeplants.msu.edu/pdf/E2973.pdf>. [Date accessed: September 26, 2007].
- Jones, G.A.; Gillett, J.L. 2005. Intercropping with sunflowers to attract beneficial insects in organic agriculture. *Florida Entomologist*. 88: 91-96.
- Landis, D.; Fiedler, A. 2005. Enhancing biological control with native plants. East Lansing, MI: Michigan State University. <http://nativeplants.msu.edu/>. [Date accessed: September 26, 2007].
- Nicholls, C.I.; Parrella, M.P.; Altieri, M.A. 2000. Reducing the abundance of leafhoppers and thrips in northern California organic vineyard through maintenance of full season floral diversity with summer cover crops. *Agricultural and Forest Entomology*. 2: 107-113.
- Pickett, C.H.; Bugg, R.L. 1998. Enhancing biological control: habitat management to promote natural enemies of agricultural pests. Berkeley, CA: University of California Press. 421 p.
- Stamps, W.T.; Linit, M.J. 1998. Plant diversity and arthropod communities: implications for temperate agroforestry. *Agroforestry Systems*. 39: 73-89.