



5.8 Windbreaks for livestock

Windbreaks located perpendicular to winter winds can protect livestock while allowing cooling summer winds to circulate in the feedlot or pasture. Protect windbreaks from grazing which can reduce windbreak effectiveness. Runoff from feedlot should be directed away from trees and should be treated (see section 1.0).

5.8 References

- Bird, P.R. 1998. Tree windbreaks and shelter benefits to pasture in temperate grazing systems. *Agroforestry Systems*. 41: 35-54.
- Dronen, S.I. 1988. Layout and design criteria for livestock windbreaks. *Agriculture, Ecosystems and Environment*. 22/23: 231-240.
- Forman, R.T.T.; Buadry, J. 1984. Hedgerows and hedgerow networks in landscape ecology. *Environmental Management*. 8: 495-510.
- Heisler, G.M.; DeWalle, D.R. 1988. Effects of windbreak structure on wind flow. *Agriculture, Ecosystems and Environment*. 22/23: 41-69.
- Loeffler, A.E.; Gordon, A.M; Gillespie, T.J. Optical porosity and windspeed reduction by coniferous windbreaks in southern Ontario. *Agroforestry System*. 17: 119-133.
- Quam, V.C.; Johnson, L.; Wight, B.; Brandle, J.R. 1994. Windbreaks for livestock production. EC 94-1766-X. Lincoln, NE: University of Nebraska Cooperative Extension. 6 p. <http://www.unl.edu/nac/morepublications/ec1766.pdf>. [Date accessed: September 26, 2007].

Read, R.A. 1957. Effect of livestock concentration on surface-soil porosity within shelterbelts. *Journal of Forestry*. 55: 529-530.

Shaw, D.L. 1988. The design and use of living snow fences in North America. *Agriculture, Ecosystems and Environment*. 22/23: 351-362.

Timm, R.M. 1988. Vertebrate pest management in windbreak systems. *Agriculture, Ecosystems and Environment*. 22/23: 555-570.