


# Where's the Beef?


Why 20 Years of Predicted Global Warming Effects on Fish Distributions Remain Unsubstantiated

Dan Isaak and Bruce Rieman (retired, sort of...)

US Forest Service - Air, Water & Aquatics Program  
Rocky Mountain Research Station  
Boise, ID 83702



# Scientific Consensus That Global Warming Would Occur for 30+ Years




Peterson et al. 2008. *Bull. Amer. Metero. Soc.* 1325-1337.





# Strong Empirical Support for Warming

## Global Air Temperature Trend




## Western U.S. Air Temperature Trends 1950 - 2009




# Regional Trends In Northwest Rivers

## Fraser River - Annual




Morrison et al. 2002

## Columbia River - Summer




Crozier et al. 2008

## Snake River, ID - Summer



Isaak et al. 2011. *Climatic Change*


## Missouri River, MT - Summer






# Temperature is Primary Control for Ectotherms Like Fish


## Temperature & metabolic rates




## Thermal Niche



## In the lab...





## & the field



# Are Species Distributions Shifting?


## Temporal distribution shifts






# Shifts in Salmon Migration Timing


## Median Spring Chinook Migration Dates at Bonneville



## Distribution of Sockeye Migrations at Lower Granite



## July Stream Temps at Bonneville





## Studies...

- Juanes et al. 2003
- Crozier et al. 2008
- Keefer et al. 2009
- Wedekind & Kung 2010
- Crozier et al. 2011
- Etc.

# Are Species Distributions Shifting?

## Spatial distribution shifts



Average distribution shift  
across taxa =  
6.1 km/decade poleward  
OR  
6.1 m/decade higher

Parmesan and Yohe. 2003.  
*Nature* 421:37-42.





# We've Predicted It for 20+ Years...


## Early brook trout climate assessments

### Effect of Climatic Warming on the Southern Margins of the Native Range of Brook Trout, *Salvelinus fontinalis*

J. Donald Meisner<sup>1</sup>

### The Role of Groundwater in the Impact of Climate Warming on Stream Salmonines

J. D. Meisner, J. S. Rosenfeld, and H. A. Regier





### Similar climate studies

- Meisner 1988, 1990
- Eaton & Schaller 1996
- Keleher & Rahel 1996
- Rahel et al. 1996
- Mohseni et al. 2003
- Flebbe et al. 2006
- Rieman et al. 2007
- Kennedy et al. 2008
- Williams et al. 2009
- Isaak et al. 2010
- Wenger et al. 2011
- Etc.

Meisner et al. 1988. *Fisheries* 13(3):2-8; Meisner 1990. *CJFAS* 47:1065-1070

# Biological Validation Doesn't Exist



Fact or Fiction?





# There's A Lot on the Line

## Climate Boogeyman



## Recreational Fisheries

Low Flows Prompt Fishing Closure On Upper Beaverhead River And Reduced Limits On Clark Canyon Reservoir

Wednesday, September 29, 2004  
Fishing


High Water  
Temperature In Grande  
Ronde Kills 239 Adult  
Spring Chinook  
Columbia Basin Bulletin,  
August 14, 2009 (PST)



## ESA Listed Species




## Land Use & Water Development






# Western Trout Climate Assessment


Fish survey database  
~10,000 sites




Historic  
Distributions





GCM



Distributions  
for IPCC A1B  
Scenarios



Species-Specific  
Habitat Response  
Curves



Wenger et al. 2011. *Proc. Nat. Acad. Sciences*

50% Reduction by  
2080




# Why Doesn't Biological Validation Exist?

We're not sampling the right places



# Why Doesn't Biological Validation Exist?

Need to sample across thermal boundaries








# What is an Isotherm?

## How Does it Apply to Streams?



Line connecting locations with equal temperatures



Plan view



Longitudinal view




Salmon River  
FLIR profile



# Key BioClimate Model Assumption:

Critical isotherm delimits population boundary






# Regional BioClimatic Model Predictions are Not Testable



Temperature isotherms mapped  
instead of fish distributions

OR



Statistically imprecise


- Bull trout lower elevation limit  $x = 1,567\text{m}$ , 95% CI = 172m
- 52 years for detectable range shift (assuming +0.2 C/decade)



Rieman et al. 2007

# Stream-Specific Predictions of Isotherm Shifts Needed for Precision

- 1) Stream temperature lapse rate ( $^{\circ}\text{C} / 100 \text{ m}$ )
- 2) Long-term stream warming rate ( $^{\circ}\text{C} / \text{decade}$ )
- 3) Stream slope (degrees)
- 4) Stream sinuosity






# A Use for High School Trigonometry!


1. Calculate vertical displacement for a given stream lapse rate and long-term warming rate.

$$\text{Displacement (a)} = \frac{\text{Warming rate}}{\text{Lapse rate}} = \frac{0.2^{\circ}\text{C/decade}}{0.4^{\circ}\text{C/100m}} = +50\text{m/decade}$$

2. Translate displacement to distance along stream of a given slope.




3. Multiply slope distance by stream sinuosity ratio in meandering streams.



# Isotherm Shift Rate Curves


Stream lapse rate = 0.4 °C / 100 m






# Isotherm Shift Rate Curves

Stream lapse rate = 0.4 °C / 100 m



# Isotherm Shift Rate Curves

Stream lapse rate = 0.8 °C / 100 m





# Mapping Climate Change "Velocity"


Long-term stream warming rate =  $0.2^{\circ}\text{C}/\text{decade}$

Stream lapse rate =  $0.4^{\circ}\text{C} / 100\text{ m}$

ISR  
(km/decade)




## Boise River Network





# Climate Vulnerability & Physiography

Trouble?




Elevational  
Refuge

Latitudinal  
Refuge






# Climate Vulnerability & Physiography



# Precise Isotherm Shift Predictions

Is it a problem?





# Precise Isotherm Shift Predictions

Is it a problem?

How much time left on the clock?

Headwater populations  
with  $\leq 10$  stream km in  
trouble by 2050

Elevation




x years until  
thermally suitable  
habitat disappears





# Biological Monitoring Implications

Longitudinal surveys to map population boundaries & establish baseline





# Measure Shift Between Surveys




TRENDS in Ecology & Evolution


Tingley & Bessinger. 2009. Detecting range shifts from historical species occurrences. *TREE* 24:625-633.

# Power Analysis for Trend Detection

How long would monitoring have to occur?



Streams differ in thermal variation & this variation partially masks climate signal that populations receive



Stream	Summer SD	Annual SD
NFK Clearwater	1.41	0.70
Fir Creek	0.82	0.51
Missouri R.	1.17	0.64
SFK Bull River	0.86	0.55
NFK Bull River	0.36	0.44
Bull River	0.82	0.58


Isaak et al. 2011.  
*Climatic Change*



# Power Curves for Isotherm Shifts


Stream lapse rate =  $0.4\text{ }^{\circ}\text{C} / 100\text{ m}$

Stream slope = 4%



# Power Curves for Isotherm Shifts

Stream lapse rate =  $0.4\text{ }^{\circ}\text{C} / 100\text{ m}$   
Stream slope = 1%



Realistic ISRs  
for 1% channels



# Empirical Evidence in the Short-Term

## Resample stream profiles from 20+ years ago



ALTITUDINAL DISTRIBUTION OF BROWN TROUT AND OTHER FISHES IN A HEADWATER TRIBUTARY OF THE SOUTH PLATTE RIVER, COLORADO

ROBERT E. VINCENT AND WILLIAM H. MILLER<sup>1</sup>

*Colorado Cooperative Fishery Unit, Colorado State University, Fort Collins, Colorado 80521*

(MS received August 9, 1968; accepted March 10, 1969)

**Fish Assemblages and Habitat Gradients in a Rocky Mountain–Great Plains Stream: Biotic Zonation and Additive Patterns of Community Change**

FRANK J. RAHEL

*Department of Zoology and Physiology, University of Wyoming  
Laramie, Wyoming 82071, USA*

WAYNE A. HUBERT

*Transactions of the American Fisheries Society 120:319–332, 1991*


Species	Site number and elevation (m)									
	1 2,234	2 2,030	3 2,015	4 1,591	5 1,559	6 1,524	7 1,510	8 1,490	9 1,470	10 1,423
	<b>Upstream species</b>									
(1) Brook trout	100	79	72	6						
(2) Brown trout		21	24	+	+	+				2
	<b>Species showing an additive pattern</b>									
(1) White sucker		4	42	34	13	26	51	35	39	
(2) Longnose dace			23	18	52	6		33	12	
(3) Longnose sucker			2	19	12	13	1	1	2	
(4) Creek chub			27	27	20	6	47	28	24	
(5) Sand shiner						1	12	+	14	
(6) Bigmouth shiner							6	+	4	
(7) Fathead minnow							8		+	
(8) Common shiner									4	
(9) Brassy minnow									+	

## DISTRIBUTION AND ABUNDANCE OF FISHES IN SAGEHEN CREEK, CALIFORNIA

RICHARD GARD, School of Forestry and Conservation, University of California, Berkeley 94720<sup>1</sup>




GLENN A. FLITTNER, Bureau of Marine Sciences, California State University, San Diego 92100

*J. Wildl. Manage.* 38(2):1974



# Broad Distributional Resurveys

Assess site extirpation/colonization frequencies relative to temperature



Beever et al. 2003; 2010



# Broad Distributional Resurveys

Assess site extirpation/colonization frequencies relative to temperature

United States  
Department of  
Agriculture  
  
Forest Service  
  
Intermountain  
Research Station  
  
General Technical  
Report INT-241  
  
February 1988


## Density and Biomass of Trout and Char in Western Streams

RELATIONSHIPS AMONG STREAM ORDER, FISH  
POPULATIONS, AND AQUATIC GEOMORPHOLOGY  
IN AN IDAHO RIVER DRAINAGE




Platts  
70's/80's

Bjornn  
1960's/70's



Fish survey database  
~10,000 sites



Wenger et al. 2011. *PNAS*



# Conclusions/Discussion

- Estimates of biological shift rates is *the "X Prize"* and critical information necessary to facilitate accurate climate risk assessments & *empower managers to make tough decisions.*
- Monitoring efforts should focus on streams with fast ISRs and low thermal variance. Detection of biological shifts will require a minimum of 20 years (but could be much longer).
- Resurveys of historical sites are needed to provide empirical evidence of biological shifts in near future.
- Headwater populations that occupy < 10 km of stream & lack upstream elevational refuges may be extirpated by 2050.
- Interesting ecological questions:
  - a) Do shift rates differ between temperature mediated boundaries where populations are allopatric or sympatric (with nonnative competitors)?
  - b) Do shift rates differ at warm (extirpation) or cold (colonization) boundaries?



The End

